

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 375 388 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication of patent specification: 02.03.94 (51) Int. Cl.⁵: A61K 7/06, A61K 7/48

(21) Application number: 89313334.8

(22) Date of filing: 20.12.89

(54) Cosmetic composition.

(30) Priority: 22.12.88 GB 8830019

(43) Date of publication of application:
27.06.90 Bulletin 90/26

(45) Publication of the grant of the patent:
02.03.94 Bulletin 94/09

(94) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI NL SE

(56) References cited:
EP-A- 0 242 967
EP-A- 0 277 428

(73) Proprietor: UNILEVER PLC
Unilever House
Blackfriars
P.O. Box 68
London EC4P 4BQ(GB)

(94) Designated Contracting States:
GB

(73) Proprietor: UNILEVER N.V.
Weena 455
NL-3013 AL Rotterdam(NL)

(94) Designated Contracting States:
BE CH DE ES FR GR IT LI NL SE AT

(72) Inventor: Davis, Michael Arthur Frank
6 Perbeck Road
Rushden
Northamptonshire NN10 9UH(GB)
Inventor: Gibson, Walter Thomas
8 Braid Court
Wellingborough
Northamptonshire NN8 3PF(GB)

(74) Representative: Ford, Michael Frederick et al
MEWBURN ELLIS
2 Cursitor Street
London EC4A 1BQ (GB)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

EP 0 375 388 B1

Description

The invention relates to cosmetic and pharmaceutical compositions for topical application to mammalian skin or hair, containing a hair growth promotor which is capable of increasing or maintaining hair growth, especially terminal hair growth on the human scalp.

BACKGROUND

The Hair Growth Cycle

It should be explained that in most mammals, hair does not grow continuously, but undergoes a cycle of activity involving alternate periods of growth and rest. The hair growth cycle can be divided into three main stages, namely:

- (i) an active stage known as anagen, during which the hair follicle penetrates deep into the dermis with the cells of the bulb dividing rapidly and differentiating to form the hair,
- (ii) a regressive stage known as catagen, which is heralded by the cessation of mitosis, and during which the follicle regresses upwards through the dermis and hair growth ceases,
- (iii) a resting stage known as telogen, in which the regressed follicle contains a small secondary germ with an underlying ball of tightly packed dermal papilla cells.

The initiation of a new anagen stage is revealed by rapid proliferation in the germ, expansion of the dermal papilla and elaboration of basement membrane components. The hair cycle is then repeated many times until, as a consequence of the onset of male pattern baldness, most of the hair follicles spend an increasing proportion of their time in the telogen stage, and the hairs produced become finer, shorter, and less visible; this is known as terminal to vellus transformation.

PRIOR ART

Alleged Baldness Cures

Although there have been many claims in the scientific literature to the promotion or maintenance of hair growth by the topical application of hair tonics and the like, with the possible exception of minoxidil, none has been shown to be sufficiently free from disadvantageous clinical side effects, whether administered topically, orally or systemically, to warrant commercial exploitation as an ethical pharmaceutical, proprietary medicine, or as a cosmetic product. Possibly, the only means which has met with partial success for growing hair on the bald or balding human head is by transplantation of hair to the bald areas. This is usually a painful operation and is not always successful. Furthermore, it is immediately apparent to the casual observer that the subject has received a hair transplant and it may take many months or even years before hair regrowth, following this operation, assumes an appearance which resembles that of the original naturally growing hair.

Among the many hair regrowth studies that have been reported in the literature, there is included the work of Bazzano as described in PCT International Publication No. WO 85/04577. This publication describes a composition which is useful for increasing the rates of hair growth on mammalian skin, for prolonging anagen and for treating various types of alopecias. The composition in question comprises a pyrimidine carbamate.

It has also been reported in US patent no. 4 139 619 to Chidsey assigned to the Upjohn Company, that a topical composition comprising minoxidil as the free base or acid addition salt thereof, or certain specified related iminopyrimidines, is useful in stimulating the conversion of vellus hair to growth as terminal hair, as well as increasing the rate of growth of terminal hair.

In spite of the apparent stimulation of hair growth or regrowth reported independently by Bazzano and Chidsey, following topical application of minoxidil or related compounds, there is general concern that systemic side-effects can result, particularly following topical application of minoxidil. Thus it is generally recognised in the medical literature that the side effects of orally administered minoxidil are very serious, and include fluid retention, tachycardia, dyspnea, gynecomastia, fatigue, nausea and cardiotoxicity. There is also evidence that certain side effects have been experienced following topical application of minoxidil.

In addition to the alleged benefits of employing the pyrimidine carbamates of Bazzano or minoxidil of Upjohn, many other hair regrowth studies have been reported in the literature.

Unilever in EP 0 277 428 report that glycosaminoglycan breakdown is an important early change in catagen, and since there is already evidence for a link between the presence of intact glycosaminoglycans

and hair growth, they suggest that prevention of proteoglycan and glycosaminoglycan breakdown may lead to earlier onset and/or prolongation of anagen. This would effectively retard hair loss and reverse baldness. Unilever also suggest that such breakdown may be prevented in a number of ways, viz by inhibiting proteoglycanase activity, by blocking cellular uptake of intact glycosaminoglycan chains, and/or by inhibiting glycosaminoglycanase activity. With particular reference to glycosaminoglycanase activity, Unilever have shown that glucaro-1,4-lactone, and related aldonolactones, inhibitors of exoglycosidases, can be topically applied to human skin to induce, maintain or increase hair growth.

We have now identified a new inhibitor of exoglycosidases, which is also believed to prevent the breakdown of glycosaminoglycan chains, and we have accordingly found that this new inhibitor will stimulate hair growth.

DEFINITION OF THE INVENTION


Accordingly, the invention provides a preserved composition suitable for topical application to mammalian skin or hair for inducing, maintaining or increasing hair growth which comprises, as the hair growth promoter, a hexosaccharic acid and/or a salt or ester thereof; the total amount of hexosaccharic acid or salt or ester thereof present in the composition being sufficient to increase hair growth in the rat, when the composition is applied topically thereto over a period of no more than 3 months, by at least 10% more than that obtainable using a control composition from which the promoter has been omitted, in accordance with the Rat Hair Growth Test.

By "preserved composition", we mean that the composition is free from viable microbial contaminants capable of resulting in microbial spoilage of the composition and/or biodegradation of the hair growth promoter.

DISCLOSURE OF THE INVENTION

The Hair Growth Promoter

The composition according to the invention comprises, as a hair growth promoter, an hexosaccharic acid or an acylated hexosaccharic acid, or a salt or ester thereof, having the structure (1).



where

- X^1 is chosen from H, alkalimetal, ammonium and substituted ammonium counter ions;
- X^2 is chosen from an alkyl or hydroxyalkyl group having from 1 to 18 carbon atoms;
- $\text{Y}^1, \text{Y}^2, \text{Y}^3$ and Y^4 are each chosen from H, an alkyl group having from 1 to 12 carbon atoms, and an acyl group having from 1 to 18 carbon atoms;
- l is an integer of from 1 to 3;
- m and n are each 0 or the integer 1 or 2; and
- $m + n$ is 1 or 2.

Examples of hexosaccharic acids, in which $\text{X}^1, \text{Y}^1, \text{Y}^2, \text{Y}^3$ and Y^4 in the above structure are -H, n is 2, and m is 0, include:

- Allosaccharic acid
- Altrosaccharic acid
- Glucosaccharic acid

Mannosaccharic acid
 Gulosaccharic acid
 Idosaccharic acid
 Galactosaccharic acid, and
 Talosaccharic acid.

Examples where X^1 is a cation, are the monovalent alkali metal cations Na^+ and K^+ .

Further examples where X^1 is a cation are substituted ammonium cations, such as diethanolammonium and triethanolammonium cations.

Examples where X^2 is an alkyl group are methyl, ethyl, n-propyl, n-butyl, n-octyl and lauryl.

Examples where Y^1 , Y^2 , Y^3 and Y^4 are alkyl groups, are methyl and ethyl.

Examples where Y^1 , Y^2 , Y^3 and Y^4 are acyl group, are acetyl and propionyl.

A particularly preferred hexosaccharic acid is glucosaccharic acid (also known as saccharic acid or glucaric acid, and hereinafter referred to as glucaric acid) having the structure (2):



A particularly stable salt of glucaric acid which is preferred, is the disodium salt.

The total amount of hexosaccharic acid, or salt or ester thereof, as the hair growth promoter, present in the composition according to the invention is sufficient to increase hair growth in the rat, the model selected for this test, when said composition is applied topically thereto by, over a period of no more than 3 months, at least 10% more than that obtainable using a control composition from which the promoter has been omitted, in accordance with the Rat Hair Growth Test to be described later.

Preferably, the amount the hair growth promoter should be sufficient to increase hair growth in the rat by at least 20%, more preferably by at least 30%, most preferably by at least 40% and ideally by at least 50%.

The sufficient amount of the promoter is from 0.0001 to 99%, preferably from 0.1 to 20% by weight of the composition.

Preservation of the Composition

The composition according to the invention is preserved in such a manner that it will enjoy an extended shelf life following manufacture and prior to sale and use. Ideally the composition will have an indefinite shelf life.

It is accordingly apparent that the hair growth promoter is particularly prone to attack by bacteria, moulds and fungi and other microbial influences, particularly at pH values near neutrality that characterise the preferred composition. The shelf-life of the composition can therefore be unacceptably short due to the biodegradation of the promoter, unless steps are taken to preserve the composition.

In order to be preserved, the composition will accordingly be free, or substantially free, from viable microbial contaminants that are capable of resulting in microbial spoilage of the composition, and/or biodegradation of the promoter prior to topical application of the composition to mammalian skin or hair. It is to be understood, however, that the invention is also concerned with compositions, as herein defined, which may contain viable but dormant micro organisms, such as bacterial spores, provided that the conditions of preservation do not result in substantial proliferation of the microorganisms prior to use of the composition.

Examples of methods that can be employed to achieve preservation of the composition, includes the following:

(i) Sterilisation

The composition according to the invention can be preserved by sterilisation to remove or kill substantially all viable microbial contaminants. This can be achieved for example by irradiation using a lethal dose of gamma rays, by heat sterilisation or by ultrafiltration using techniques that are well established in the pharmaceutical industry.

(ii) Extremes of pH value

The composition according to the invention can alternatively be preserved by adjusting its pH to values that are either too low (eg. pH <2) or too high (eg. pH >12) to permit significant proliferation of microbial contaminants. The pH of the composition can accordingly be adjusted to desired high or low values by addition of an alkali or acid as a pH adjustant.

(iii) Chemical Preservative

The composition according to the invention can also be preserved by including in it a chemical preservative which functions to prevent the growth of or kill bacteria, fungi or other microorganisms.

Examples of chemical preservatives include ethanol, benzoic acid, sodium benzoate, sorbic acid, potassium sorbate, sodium propionate and the methyl, ethyl, propyl and butyl esters of p-hydroxybenzoic acid. The amount of chemical preservative that can be incorporated in the composition according to the invention will generally be from 0.05 to 25%, the amount chosen being sufficient to arrest microbial proliferation.

(iv) Water activity depressants

The compositions according to the invention can also be preserved by the inclusion of a water activity depressant such as glycerol, propylene glycol, sorbitol, sugars and salts, for examples alkali metal halides, sulphates and carboxylates. When employing a water activity depressant, sufficient should be incorporated in the composition according to the invention to reduce the water activity (a_w) from 1 to <0.9, preferably to <0.85 and most preferably < 0.8, the lowest of these values being that at which yeasts, moulds and fungi will not proliferate.

OTHER INGREDIENTSVehicle

The composition according to the invention also optionally comprises a solid, semi-solid or liquid cosmetically and/or physiologically acceptable vehicle, to enable the promoter to be conveyed to the skin at an appropriate dilution. The nature of the vehicle will depend upon the method chosen for topical administration of the composition. The vehicle can itself be inert or it can possess physiological or pharmaceutical benefits of its own.

The selection of a vehicle for this purpose presents a wide range of possibilities depending on the required product form of the composition. Suitable vehicles can be classified as described hereinafter.

It should be explained that vehicles are substances which can act as diluents, dispersants, or solvents for the inhibitor which therefore ensure that it can be applied to and distributed evenly over the hair and/or scalp at an appropriate concentration. The vehicle is preferably one which can aid penetration of the promoter into the skin to reach the immediate environment of the hair follicle. Compositions according to this invention can include water as a vehicle, and/or at least one cosmetically acceptable vehicle other than water.

Vehicles other than water that can be used in compositions according to the invention can include solids or liquids such as emollients, solvents, humectants, thickeners and powders, some of which can also be employed to preserve the composition.

Examples of vehicles, which can be used singly or as mixtures of one or more vehicles, are as follows:

Emollients, such as stearyl alcohol, glyceryl monoricinoleate, glyceryl monostearate, propane-1,2-diol, butane-1,3-diol, mink oil, cetyl alcohol, isopropyl isostearate, stearic acid, isobutyl palmitate, isocetyl stearate, oleyl alcohol, isopropyl laurate, hexyl laurate, decyl oleate, octadecan-2-ol, isocetyl alcohol, cetyl palmitate, dimethylpolysiloxane, di-n-butyl sebacate, isopropyl myristate, isopropyl palmitate, isopropyl stearate, butyl

stearate, polythylene glycol, triethylene glycol, lanolin, sesame oil, coconut oil, arachis oil, castor oil, acetylated lanolin alcohols, petroleum, mineral oil, butyl myristate, isostearic acid, palmitic acid, isopropyl linoleate, lauryl lactate, myristyl lactate, decyl oleate, myristyl myristate;

Propellants, such as trichlorofluoromethane, dichlorodifluoromethane, dichlorotetrafluoroethane, monochlorodifluoromethane, trichlorotrifluoroethane, propane, butane, isobutane, dimethyl ether, carbon dioxide, nitrous oxide;

Solvents, such as ethyl alcohol, methylene chloride, isopropanol, castor oil, ethylene glycol monoethyl ether, diethylene glycol monobutyl ether, diethylene glycol monoethyl ether, dimethyl sulphoxide, dimethyl formamide, tetrahydrofuran;

Humectants, such as glycerin, sorbitol, sodium 2-pyrrolidone-5-carboxylate, soluble collagen, dibutyl phthalate, gelatin;

Powders, such as chalk, talc, fullers earth, kaolin, starch, gums, colloidal silicon dioxide, sodium polyacrylate, tetra alkyl and/or trialkyl aryl ammonium smectites, chemically modified magnesium aluminium silicate, organically modified montmorillonite clay, hydrated aluminium silicate, fumed silica, carboxyvinyl polymer, sodium carboxymethyl cellulose, ethylene glycol monostearate.

The amount of vehicle in the composition, including water if present, should preferably be sufficient to carry at least a portion of the promoter to the skin in an amount which is sufficient effectively to enhance hair growth. The amount of the vehicle can comprise the balance of the composition, particularly where little or no other ingredients are present in the composition. Accordingly, the vehicle or vehicles can comprise from 1 to 99.99%, preferably from 50 to 99.5% and ideally from 90 to 99% by weight of the composition.

Perfume

The composition according to the invention can also optionally comprise a perfume in an amount sufficient to make the composition acceptable to the consumer and pleasant to use. Usually, the perfume will form from 0.01 to 10% by weight of the composition.

Activity Enhancer

The composition according to the invention can also optionally comprise an activity enhancer.

The activity enhancer can be chosen from a wide variety of molecules which can function in different ways to enhance the effects of the hair growth promoter. Particular classes of activity enhancers include other hair growth stimulants, penetration enhancers, cationic polymers and liposomes, whose presence can further improve the delivery of the promoter through the stratum corneum to its site of action in the immediate environment of the hair follicle.

Some activity enhancers can also function as vehicles for the hair growth promoter.

(a) Other Hair Growth Stimulants

i. Examples of other substances which themselves possess the ability to stimulate or increase hair growth include;

Benzalkonium chloride

Benzethonium chloride

Phenol

Estradiol

Diphenhydramine hydrochloride

Chlorpheniramine maleate

Chlorophyllin derivatives

Cholesterol

Salicylic acid

Methionine

Cystine

Red pepper tincture

Benzyl nicotinate

dl-Menthol

Peppermint oil


Calcium pantothenate

Panthenol

Castor oil
Hinokitiol
Prednisolone
Resorcinol

5 ii. Further substances which themselves possess the ability to increase the rate of terminal hair growth include:

α -1,4 esterified disaccharides described by Choay S.A. in EP-A-O 064 012, having the structure (3):



20 where


Z represents a functional nitrogen group, such as an azide or a group having the structure -NHB, in which B represents -H or a functional group such as acetyl or sulphate as a salt with an organic or mineral cation;

25 M represents -H or SO_3M_1 , where M_1 is an organic or metallic cation, particularly an alkali metal; or an acetyl group;


R represents a C_1 to C_4 alkyl radical, especially methyl; or an aryl radical;

A represents a functional group such as an acid or $-\text{COOR}_1$, where R_1 represents -H or a C_1 to C_4 alkyl radical, especially methyl; or a metal, especially an alkali metal;

30 esterified oligosaccharides as described by Unilever in EP-A-O 211 610, including at least one esterified disaccharide unit consisting of a uronic acid residue having the structure (4a):




and a hexosamine residue having the structure: (5)



where

55 R' is -H, C_3 to C_{10} alkyl or



R' is -H, C₁ to C₄ alkyl, -CO(CH₂)_mCH₃, -SO₃M,

R'' is -H, CO(CH₂)_mCH₃, or -SO₃M,

M is -H, or a metallic or organic cation

n is 0 or an integer of from 1 to 7, and

m is 0 or the integer 1 or 2;

the groups designated R' being the same or different, one R' group from each pyranose ring structure being linked by a glycosidic linkage having the configuration α-1,3, α-1,4, β-1,3 or β-1,4; and the -COOR', -CH₂OR'

and -OR' groups being of either configuration with respect to the pyranose rings;

iii. Minoxidil and certain derivatives thereof as described by the Upjohn Co. in US 3 461 461,

iv. Minoxidil glucuronides, as described by Unilever in EP-O 242 967,


v. Minoxidil sulphates, as described by The Upjohn Co. in WO 86/04231,

vi. Ethylenediaminetetraacetic acid or salts thereof, as described by Redken Laboratories, Inc. in US 4 814 351.

vii. Aldonolactones, such as D-glucaro-1,4-lactone, as described by Unilever in EP-0 277 428.


viii. Direct proteoglycanase inhibitors, such as 1,10-phenanthroline, as described by Unilever in EP-0 277 428.

ix. Glycosaminoglycanase inhibitors, as described by Unilever in EP-0 277 428, such as aldonolactones and esterified aldonolactones having the structure (5):



where

A¹ and A² are -H, -CH₃,



B is OD'' or a lactone linkage to position 1 or 6, or -NHCOCH₃ and where

D is -H or C₂ to C₈ alkyl,

D' is the remainder of the molecule joined through another C atom at positions 2 to 5 to form a lactone,

D'' is -H or C₂ (ie acetyl) to C₄ acyl of either configuration with respect to the backbone of this

molecule;

preferred examples of which include:

L-Galactono-1,4-lactone

L-Arabin-1,5-lactone

5 D-Fucono-1,5-lactone

D-Glucono-1,4-lactone

D-Glucurono-6,3-lactone

Galactaric acid lactone

2-Acetamido-2-deoxygluconolactone

10 2-Acetamido-2-deoxygalactono-lactone


D-Glucono-1,4:6,3-dilactone

L-Idaro-1,4-lactone

2,3,5-Tri-O-acetyl-D-glucono-1,4-lactone

2,5-Di-O-acetyl-D-glucono-1,4:6,3-dilactone.

15 x. Glycosaminoglycanase inhibitors, as described by Unilever in EP 0 277 428, such as monosaccharides and esterified monosaccharides having the structure (6):



where

A is -OG or -NHCOCH₃

G is -H, -SO₃M⁺, C₂ (ie acetyl) to C₄ acyl

40 G' is -H or -OG

M⁺ is -H or a metal cation

wherein the functional groups can be in either configuration with respect to the backbone of the above molecule;

preferred examples of which include:

45 N-Acetylglucosamine


N-Acetylgalactosamine

D-Galactosamine

D-Glucosamine-3-sulphate

N-Acetylmannosamine.

50 xi. Glycosaminoglycan chain cellular uptake inhibitors, as described by Unilever in EP 0 277 428, such as hexuronic acid and esters thereof which may be represented by the generic structure (7):



where


G is -H, -SO₃M⁺, C₂ (ie acetyl) to C₄ acyl;

D is -H or C₂ to C₈ alkyl

M⁺ is -H or a metal cation;

wherein the functional groups can be in either configuration with respect to the backbone of the above molecule;

xii. Chemical inhibitors of glycosidase activity, as described by Unilever in EP 0 334 586, chosen from lactams having the structure (8):



where

A³ and A⁴ are -H, -CH₃,



-CH₂OT
or



5

A³ and A⁴ being the same or different, and at least one of which being the group:

10



in a lactam ring;

and where Q is -OT', -NHT' or a lactam linkage to A³ or A⁴;

15

the Q groups being the same or different, and at least one of which is involved in a lactam linkage;

and where T is the same or different and is chosen from

-H, -C_pH_{2p+1} or a metal ion,

T' is -H or -COC_pH_{2p+1}, and

p is an integer of from 1 to 22;

20

provided that:

where any of the Q groups is

-OT' or -NHT',

then that group or groups can be of either stereochemical configuration with respect to the plane of the ring,

25

preferred examples of which include:

D-glucaro-1,5-lactam L-Galactono-1,4-lactam,

L-Arabino-1,5-lactam, D-Fucono-1,5-lactam,

D-Glucaro-1,4-lactam, D-Glucurono-6,3-lactam,

1,2,5-tri-O-acetyl-D-glucurono-6,3-lactam

30

2-Acetamido-2-deoxygluconolactam,

2-Acetamido-2-deoxygalactonolactam,

D-Glucaro-1,4:6,3-dilactam, L-Idaro-1,4-lactam,

2,3,5-Tri-O-acetyl-D-glucaro-1,4-lactam,


2,5-Di-O-acetyl-D-Glucaro-1,4:6,3-dilactam,

35

D-glucaro-1,5-lactam ethyl ester;

xiii. Chemical activators of protein kinase C enzymes, as described by Unilever in EP 0 334 585 chosen from diacylglycerols having the structure (9):

40




(9)

45

50

where X and X' are the same or different and is represented by the grouping:



55

(CH=CH)_b] CH₃


where x is 0 or an integer of from 1 to 28, and y is 0 or an integer of from 1 to 5;

the X and X' groups being of either stereochemical configuration with respect to the carbon backbone of the glycerol molecule;


preferred examples of which include:

1,2-Dibutanoyl-rac-glycerol
 1,2-Dihexanoyl-sn-glycerol
 1,2-Dioctanoyl-rac-glycerol
 1,2-Dioctanoyl-sn-glycerol
 1,2-Didecanoyl-rac-glycerol
 1-Oleoyl-2-acetyl-rac-glycerol
 1-Oleoyl-2-acetyl-sn-glycerol
 1-Stearoyl-2-arachidonoyl-sn-glycerol
 1,2-Distearoyl-rac-glycerol
 1,2-Dipentadecanoyl-sn-glycerol
 1,2-dipentadecanoyl-rac-glycerol
 1,2-Dipalmitoyl-rac-glycerol
 1,2-Dipalmitoyl-sn-glycerol
 1,2-Diseptadecanoyl-rac-glycerol
 1,2-Dioleoyl-sn-glycerol
 1,2-Dioleoyl-rac-glycerol
 1,2-Diarachidonoyl-sn-glycerol
 1,2-Dieicosanoyl-sn-glycerol
 1,2-Didoecosanoyl-rac-glycerol, and
 1,2-Dioctaeicosanoyl-sn-glycerol.


xiv. Glycosaminoglycanase inhibitors, as described by Unilever in EPA 89306278.6, chosen from aldonomonolactone or alduronomonolactone derivatives having the structure (10):



where A^5 is



A^6 is



or CH_2OR^6

B^1 , B^2 , B^3 and B^4 are each chosen from is OR^5 , NHR^6 , NHR^7 or a lactone linkage to position 1 or 6, and/or an ether linkage to Q^1 ;

said substituents B being the same or different, and being in either configuration, with respect to the backbone of the above structure, on positions C^2 to C^5 not involved in a lactone ring;

and where

R^4 is -H, C_1 to C_{20} alkyl, a metal cation, NH_4^+ or an alkanolamine cation;

R^5 is the remainder of the molecule joined through another C atom at positions 2 to 5 to form a lactone;

R^6 is -H, $-\text{CH}_3$, benzyl or C_2 to C_6 acyl; R^7 is -H, $-\text{CH}_3$, benzyl or C_3 to C_6 acyl;

Q^1 is the remainder of the molecule joined through an ether linkage to either C^4 or C^5 , forming either a pyranose or furanose ring;

provided that, when A^5 is



then A^6 is



provided also that, when A^6 is CH_2OH , then one or more of the B substituents is $-\text{CH}_3$, C_2 to C_4 acyl or NHR^7 ;

provided also that, when A^5 is



and all B^1 , B^2 , B^3 and B^4 substituents are -OH, then A^6 is



or CH_2OR^6 , and R^4 is C_1 or C_3 to C_{20} alkyl;

preferred examples of which aldonomonolactone derivatives include:

6-acetyl-galactono-1,4-lactone

6-propionyl-galactono-1,4-lactone

6-butyryl-galactono-1,4-lactone

2-propionamido-2-deoxygluconolactone

2-butyramido-2-deoxygluconolactone

2-propionamido-2-deoxygalactonolactone

2-butyramido-2-deoxygalactonolactone

6-propionyl-2-acetamido-2-deoxygluconolactone

diacetyl-6-propionyl-2-acetamido-2-deoxygluconolactone

6-butyryl-2-acetamido-2-deoxygalactonolactone

diacetyl-6-butyryl-2-acetamido-2-deoxygalactonolactone

2,3,5,6-tetraacetyl-galactono-1,4-lactone

2,3,5-triacetyl-6-propionylgalactono-1,4-lactone

triacetyl-2-propionamido-2-deoxygalactonolactone

triacetyl-2-butyramido-2-deoxygluconolactone

6-methyl-glucaro-1,4-lactone

2,3,5,6-tetramethyl-glucaro-1,4-lactone

6-methyl-2,3,5-triacetylglucaro-1,4-lactone


6-methyl-3-methyl-glucaro-1,4-lactone, and

6-methyl-3-acetyl-glucaro-1,4-lactone;

and a preferred example of which alduronomonolactone derivative is:

1,2,5-triacetyl-glucurono-6,3-lactone.

xv. Glycosaminoglycanase inhibitors, as described by Unilever in EPA 89306278.6, chosen from acylated monosaccharides having the structure (11):




where

A^7 is -OY or -NHR⁸

B^5 and B^6 are each chosen from is -OY, or an ether linkage to D^1 ,

D^1 is



where X^2 is an ether linkage either to C^4 or C^5 forming a pyranose or furanose ring;

Y is -H, -SO₃M, C₂ to C₄ acyl or C₁ to C₁₈ alkyl;

said substituents A^7 , B^5 , B^6 and -OY being the same or different, and being in either configuration, with respect to backbone of the above structure;

and where

Z^1 is -H or -OY

R^8 is -H, -SO₃M² or C₃ or C₄ acyl,

M^2 is -H, a metal cation, NH₄⁺, or an alkanolamine cation;

provided that, when R^8 is -H, then 1 or more of Y is chosen from -SO₃M² or C₂ to C₄ acyl; and mixtures thereof.

Preferred examples of which acylated monosaccharides include:

2-propionamido-2-deoxyglucose


1,3,4,6-tetraacetyl-2-propionamido-2-deoxyglucose

2-butyramido-2-deoxygalactose

1,3,4,6-tetraacetyl-2-butyramido-2-deoxygalactose

2-sulphamido-2-deoxygalactose
 2-sulphamido-2-deoxyglucose
 2-butyramido-2-deoxymannose
 1,3,4,6-tetraacetyl-2-butyramido-2-deoxymannose
 2-butyramido-2-deoxyglucose, and
 1,3,4,6-tetraacetyl-2-butyramido-2-deoxyglucose.

xvi. Esters of pyroglutamic acid, as described by Lever Brothers Company in US patent No. 4 774 255, having the structure (12):




(12)

where

R^1 is C_1 to C_{30} alkyl, or



and where R^2 and R^3 are the same or different and are each represented by H or the grouping (13):



where

u is zero or 1

v is zero, or the integer 1 or 2,

w is zero, or an integer of from 1 to 21

x is zero, or an integer of from 1 to 4,

y is zero, or the integer 1 or 2,

z is zero, or an integer of from 1 to 4, and

$u + v + w + x + y + z$ is an integer of from 1 to 22;

provided that when the subgrouping $(CH=CH)$ is present, then the total number of carbon atoms in said grouping is from 10 to 22.

Examples of suitable esters of pyroglutamic acid where R^1 in structure (12) is C_1 to C_{30} alkyl are:

pyroglutamic acid methyl ester

pyroglutamic acid ethyl ester

pyroglutamic acid n-propyl ester

pyroglutamic acid n-butyl ester

pyroglutamic acid n-hexyl ester

pyroglutamic acid n-heptyl ester

pyroglutamic acid n-octyl ester

pyroglutamic acid n-nonyl ester

pyroglutamic acid n-decyl ester

pyroglutamic acid n-undecyl ester

pyroglutamic acid n-dodecyl ester

pyroglutamic acid n-tridecyl ester

pyroglutamic acid n-tetradecyl ester

pyroglutamic acid n-hexadecyl ester

pyroglutamic acid n-octadecyl ester

pyroglutamic acid n-eicosyl ester

pyroglutamic acid iso-propyl ester

pyroglutamic acid 2-methylhexyl ester

pyroglutamic acid 2-ethylhexyl ester
 pyroglutamic acid 3,7-dimethyloctyl ester
 pyroglutamic acid 2-hexyldecyl ester
 pyroglutamic acid 2-octyldodecyl ester
 5 pyroglutamic acid 2,4,4-trimethyl-1-pentane ester
 pyroglutamic acid methyloctyl ester

Particularly preferred esters of this group are those where R¹ in structure (12) is C₁ to C₁₄ alkyl, (linear or branched), especially C₁ to C₆ (linear or branched).

Further examples of preferred esters of pyroglutamic acid, where R¹ in structure (12) is



15 are those where R² and/or R³ having the structure shown for grouping (13), include straight and branched chain, saturated or unsaturated aliphatic groups having from 1 to 22 carbon atoms, such as the alkyl groups:

20 methyl
 ethyl
 propyl
 iso-propyl
 butyl
 iso-butyl
 25 n-valeryl
 iso-valeryl
 n-caproyl
 n-heptyl
 n-caprylyl
 30 n-capryl
 lauryl
 myristyl
 palmityl
 stearyl, and
 35 arachidyl.

and the C₁₀₋₂₂ alkenyl groups:

linoleyl
 linolenyl
 γ-linolenyl
 40 arachidonyl, and
 columbinylyl.

Further examples of the grouping (13) also include hydroxyalkyl groups having from 1 to 22 carbon atoms, such as:

45 hydroxymethyl
 2-hydroxyethyl
 2-hydroxy-n-propyl
 3-hydroxy-n-propyl
 2-hydroxy-n-butyl
 3-hydroxy-n-butyl
 50 4-hydroxy-n-butyl
 5-hydroxy-n-valeryl
 6-hydroxy-n-caproyl
 2,3-dihydroxy-n-propyl
 2,3-dihydroxy-n-butyl
 55 12-hydroxystearyl.

Further specific examples of esters of pyroglutamic acid which are particularly suited for use as other hair growth stimulants are:

2-[pyroglutamoyloxy]-propionic acid

methyl-2-[pyroglutamoyloxy]-acetate
 ethyl-2-[pyroglutamoyloxy]-n-propionate
 ethyl-2-[pyroglutamoyloxy]-n-butyrate
 ethyl-2-[pyroglutamoyloxy]-iso-butyrate
 5 ethyl-2-[pyroglutamoyloxy]-n-valerate
 ethyl-2-[pyroglutamoyloxy]-n-caproate
 ethyl-2-[pyroglutamoyloxy]-n-heptylate
 ethyl-2-[pyroglutamoyloxy]-n-caprylate
 ethyl-2-[pyroglutamoyloxy]-n-pelargonate
 10 ethyl-2-[pyroglutamoyloxy]-3-hydroxybutyrate
 iso-propyl-2-[pyroglutamoyloxy]-n-propionate
 iso-propyl-2-[pyroglutamoyloxy]-n-caprylate
 n-propyl-2-[pyroglutamoyloxy]-n-propionate
 n-propyl-2-[pyroglutamoyloxy]-n-caprylate
 15 stearyl-2-[pyroglutamoyloxy]-n-propionate
 12-hydroxystearyl-2-[pyroglutamoyloxy]-n-propionate
 stearyl-2-[pyroglutamoyloxy]-n-stearate
 palmityl-2-[pyroglutamoyloxy]-n-propionate
 linoleyl-2-[pyroglutamoyloxy]-n-propionate
 20 linoleyl-2-[pyroglutamoyloxy]-n-caprylate
 lauryl-2-[pyroglutamoyloxy]-n-caprylate
 stearyl-2-[pyroglutamoyloxy]-n-caprylate
 glyceryl mono(2-[pyroglutamoyloxy]-n-propionate)
 glyceryl mono(2-[pyroglutamoyloxy]-n-caprylate), and
 25 glyceryl di(2-[pyroglutamoyloxy]-n-propionate).

(b) Penetration Enhancers

As has been stated earlier, the presence of a penetration enhancer can potentiate the benefit of the hair
 30 growth promoter, by improving its delivery through the stratum corneum to its site of action in the
 immediate environment of the hair follicle close to the dermal papilla.

The penetration enhancer can accordingly function in a variety of ways. It can for example, improve the
 distribution of the hair growth promoter on the skin surface or, it can increase its partition into the skin from
 the composition when applied topically, so aiding its passage to its site of action. Other mechanisms
 35 enhancing the benefit of the promoter may also be involved.

Examples of penetration enhancers include:

2-methyl propan-2-ol
 Propan-2-ol
 Ethyl-2-hydroxypropanoate
 40 Hexan-2,5-diol
 POE(2) ethyl ether
 Di(2-hydroxypropyl) ether
 Pentan-2,4-diol
 Acetone
 45 POE(2) methyl ether
 2-hydroxypropionic acid
 2-hydroxyoctanoic acid
 Propan-1-ol
 1,4 Dioxane
 50 Tetrahydrofuran
 Butan-1,4-diol
 Propylene glycol dipelargonate
 Polyoxypropylene 15 stearyl ether
 Octyl alcohol
 55 POE ester of oleyl alcohol
 Oleyl alcohol
 Lauryl alcohol
 Dioctyl adipate

- Dicapryl adipate
- Diisopropyl adipate
- Diisopropyl sebacate
- Dibutyl sebacate
- 5 Diethyl sebacate
- Dimethyl sebacate
- Dioctyl sebacate
- Dibutyl suberate
- Dioctyl azelate
- 10 Debenzyl sebacate
- Dibutyl phthalate
- Dibutyl azelate
- Ethyl myristate
- Dimethyl azelate
- 15 Butyl myristate
- Dibutyl succinate
- Didecyl phthalate
- Decyl oleate
- Ethyl caproate
- 20 Ethyl salicylate
- Isopropyl palmitate
- Ethyl laurate
- 2-ethyl-hexyl pelargonate
- Isopropyl isostearate
- 25 Butyl laurate
- Benzyl benzoate
- Butyl benzoate
- Hexyl laurate
- Ethyl caprate
- 30 Ethyl caprylate
- Butyl stearate
- Benzyl salicylate
- 2-hydroxypropanoic acid
- 2-hydroxyoctanoic acid,
- 35 Further examples of penetration enhancers include:-
- Dimethyl sulphoxide
- N,N-Dimethyl acetamide
- N,N-Dimethyl formamide
- 2-Pyrrolidone
- 40 1-Methyl-2-pyrrolidone
- 5-Methyl-2-pyrrolidone
- 1,5-Dimethyl-2-pyrrolidone
- 1-Ethyl-2-pyrrolidone
- Phosphine oxides
- 45 Sugar esters
- Tetrahydrofurfural alcohol
- Urea
- Diethyl-m-toluamide, and
- 1-Dodecylazacycloheptan-2-one
- 50 Further examples of penetration enhancers include surface active agents, preferred examples of which include:
- (i) Anionic surface active agents, such as metallic or alkanolamine salts of fatty acids for example sodium laurate and triethanolamine oleate;
- alkyl benzene sulphonates, for example triethanolamine dodecyl benzene sulphonate;
- 55 alkyl sulphates, for example sodium lauryl sulphate;
- alkyl ether sulphates, for example sodium lauryl ether sulphate [2 to 8 EO];
- sulphosuccinates, for example sodium dioctyl sulphosuccinate;
- monoglyceride sulphates, for example sodium glyceryl monostearate monosulphate;

isethionates, for example sodium isethionate;
 methyl taurides, for example Igepon TM T;
 acylsarcosinates, for example sodium myristyl sarcosinate;
 acyl peptides, for example Maypons TM and Lamepons TM ;

5 acyl lactylates,

polyalkoxylated ether glycolates, for example trideceth-7 carboxylic acid;
 phosphates, for example sodium dilauryl phosphate.

(ii) Cationic surface active agents, such as amine salts, for example sapamin hydrochloride;
 quaternary ammonium salts, for example Quaternium 5, Quaternium 31 and Quaternium 18;

10 (iii) Amphoteric surface active agents, such as imidazol compounds, for example Miranol TM ;
 N-alkyl amino acids, such as sodium cocaminopropionate and asparagine derivatives;

betaines, for example cocoamidopropylbetaine

(iv) Nonionic surface active agents, such as fatty acid alkanolamides, for example oleic ethanolamide;
 esters of polyalcohols, for example Span;

15 polyglycerol esters, for example that esterified with C₁₂₋₁₈ fatty acids and one or several OH groups;
 polyalkoxylated derivatives, for example polyoxy:polyoxyethylene stearate, and octylphenoxy polyethox-
 yethanol (TRITON TM X-100 available from Rohm & Haas, or e.g. CIRRASOL TM available from I.C.I.)

ethers, for example polyoxyethylene lauryl ether;

20 ester ethers, for example Tween TM ;

amine oxides, for example coconut and dodecyl dimethyl amine oxides.

Mixtures of two or more of the above surface active agents can be employed in the composition according to the invention.

(c) cationic polymers chosen from:

Guar Hydroxypropyltrimonium chloride

25 Quaternium-19

Quaternium-23

Quaternium-40

Quaternium-57

Poly(dipropyldiallylammonium chloride)

30 Poly(methyl- β -propaniodiallylammonium chloride)

Poly(diallylpiperidinium chloride)

Poly(vinyl pyridinium chloride)

Quaternised poly (vinyl alcohol)

Quaternised poly (dimethylaminoethylmethacrylate); and

35 mixtures thereof

The amount of activity enhancer, when employed in accordance with the invention, will normally be from 0.1 to 50%, preferably from 0.5 to 25% and most preferably from 0.5 to 10% by weight of the composition.

40 Other hair growth promoter adjuncts

There is evidence of an inflammatory component in the development of male pattern baldness and for this reason it is advantageous to include in the compositions according to the invention agents which will suppress the inflammatory process. Such agents, which include steroidal and non-steroidal anti-inflam-
 45 matory compounds (e. g. hydrocortisone and ibuprofen respectively), further enhance the hair growth promoting benefits of the hexosaccharic acid and their derivatives as herein defined, and compositions containing them, in accordance with the invention.

The composition according to the invention can also contain adjuncts other than those already mentioned, depending on the form of the intended product. It is, for example, possible to include
 50 antiseptics, antioxidants, emulsifiers, sequestrants, such as ethylene diamine, tetracetic acid and colouring agents, which can improve the stability and consumer appeal of the composition.

The composition according to the invention can also be employed as a vehicle for a wide variety of cosmetically or pharmaceutically active ingredients, particularly ingredients which have some beneficial effect other than the promotion of hair growth when applied to the skin.

55

Process

The invention also provides a process for the preparation of a composition suitable for promoting hair growth which comprises mixing a hexosaccharic acid, or a salt or ester thereof, as a hair growth promoter, with a chemical preservative, or preserving it by other means, as herein before defined, to provide a composition according to the invention, in which the promoter forms from 0.0001 to 99% by weight of the composition.

Product Form and Container

The compositions of the invention can be formulated as liquids, for example as a lotion, shampoo, milk, cream or mousse for use in conjunction with an applicator such as a roll-ball applicator, or a spray device such as an aerosol can containing propellant, or a container fitted with a pump to dispense the liquid product.

Alternatively, the compositions of the invention can be solid or semi-solid, for example sticks, creams or gels, for use in conjunction with a suitable applicator or simply a tube, bottle or lidded jar, or as a liquid-impregnated fabric, such as a tissue wipe.

The invention accordingly also provides a closed container containing a composition as herein defined.

Use of the Hair Growth Promoter for Inducing, Maintaining or Increasing Hair Growth

The invention also provides for the use of the promoter, as herein defined, and compositions containing it for topical application to mammalian skin or hair for inducing, maintaining or increasing hair growth.

The compositions according to the invention are primarily intended for topical application to the scalp of the human subject, particularly where the head is already bald or balding, in order to promote the regrowth of terminal hair. The compositions can also be applied profilactically to the hair and hence the scalp to reduce or prevent the onset of baldness.

The amount of the composition and the frequency of application to the hair and/or scalp can vary widely, depending on personal needs, but it is suggested as an example that topical application of from 0.1 to 5g daily containing from 0.00001 to 1g of a selected hair growth promoter over the period of at least six months will in most cases result in an improvement in hair growth.

EVALUATION OF EFFICACY OF HAIR GROWTH PROMOTERS USING THE RAT MODELThe Rat Hair Growth Test

The effect of compounds on hair growth was assessed using male albino Wistar rats as an animal model. The rats were chosen from as few litters as possible and were each approximately 42 days of age at the start of the test. Each rat was housed individually to prevent licking.

In each comparison, 10 rats were used in each group and hair growth was assessed as follows:

A small patch of normal skin (4cm x 4cm) on the upper back of each rat was clipped at the start and 0.3 ml of a hair growth stimulant composition (or a control) applied topically twice daily and once on Saturdays and Sundays to each clipped area. The concentration of test compound in the composition was chosen from 0.01 to 20% w/v.

It is to be understood that the potency of each hair growth promoter in terms of its ability to induce, maintain or increase hair growth is unlikely to be uniform, some being more potent than others, and therefore the concentration of any promoter chosen for thorough evaluation must be carefully selected after preliminary testing to determine its potential as a hair growth promoter. In any case, this concentration will lie within the range of from 0.01 to 20% w/v as stipulated above.

Hair was clipped from the area of the patch twice weekly, collected and weighed at each time point over a standard period of 3 months, and cumulative hair weight calculated. From these data, it was possible to estimate the effect of a promoter as a hair growth stimulant (test compound) on the amount and duration of hair growth during the experiment. A positive response, i.e. an increase of at least 10% by weight of hair after 3 months treatment, compared with a control indicates the potential of the test compound to prevent hair loss and/or reverse baldness in human subjects.

Accordingly, when the hair growth promoter is assessed compound by the Rat Hair Growth Test, an increase of at least 10% by weight of hair after 3 months treatment will be obtained. Usually, the 10% by weight minimum value will be attained well before the end of this 3 months period.

Validation of rat model for hair growth using Minoxidil

The rat model was validated by showing that twice daily topical application of a known promoter of human hair growth, namely 2% (w/v) minoxidil in a vehicle of 70% ethanol, 20% water and 10% propylene glycol, caused a significant increase of 25% in hair growth as shown below:

Table 1

Treatment	Mean Cumulative Hair weight (mg) \pm sd, after 52 days	Significance Level (vs vehicle)
2% minoxidil	786.2 \pm 94.8	p = 0.002*
Vehicle (control)	628.3 \pm 90.0	

* statistically significant

Measurement of hair growth following topical application of the disodium salt of glucaric acid

Topical treatment with a composition according to the invention was found to stimulate hair growth. In this example, the effect of topical application of the disodium salt of glucaric acid is shown. The test solution in this experiment contained 10, 15 or 20% (w/v) of the glucaric acid disodium salt in the form of a solution in a vehicle composed of 20% (v/v) ethanol, 0.1% (w/v) TRITON x-100 and 1% (w/v) citric acid adjusted to pH7 with sodium hydroxide. The control solution contained no glucaric acid disodium salt. Test or control solutions (0.3 ml) were applied topically twice daily and once on Saturdays and Sundays to the clipped area. The hair growth results are shown in Table 2.

Table 2

Treatment	Mean Cumulative Hair weight (mg) \pm sd, after 57 days	Significance Level (vs vehicle)
10% Glucaric acid	589 \pm 90.64	NS
Vehicle (Control)	563.10 \pm 81.52	(P = 0.5)
15% Glucaric acid	646 \pm 101.79	S
Vehicle (Control)	517.50 \pm 66.36	(P = 0.004)
20% Glucaric acid	998.27 \pm 142.83	S
Vehicle (Control)	763.75 \pm 138.24	(P = 0.002)
S statistically significant		
NS not statistically significant		

In addition to demonstrating a statistically significant stimulation of hair growth (up to 30% increase) as shown in Table 2, the glucaric acid disodium salt has consistently been found to advance anagen, thus reducing the amount of time spent in the resting stage of hair cycle.

Examples

The invention is illustrated by the following examples, each of which includes a hair growth promoter. Each formulation is preserved, either by irradiation or by the addition of a chemical preservative.

Furthermore, each of the formulations which features in these examples of compositions according to the invention, is capable of increasing hair growth in the rat, when applied thereto, by at least 10% more than that obtainable using a control composition from which the hair growth promoter has been omitted. It will be appreciated, however, that for human use, those of the following formulations which are "rinse off" products such as hair shampoo, will generally require repeated application to the hair or scalp before significant hair growth is observed.

Example 1

This Example illustrates a lotion according to the invention which is suitable for topical application to the scalp in order to promote hair growth.

The lotion has the following formulation:

	% w/v
Glucaric acid, methyl ester	5
ethanol	45
perfume	q.s.
water	to 100

Example 2

This Example illustrates a hair tonic which is suitable for application to hair or scalp.
The hair tonic has the following formulation:

	% w/v
disodium glucarate	15
ethanol	20
perfume	q.s.
water	to 100

Example 3

This Example also illustrates a lotion which is suitable for topical application to the scalp.
The lotion has the following formulation:

	% w/v
glucaric acid	15
propan-2-ol	10
ethanol	20
triethanolamine	q.s.*
perfume	to 100

*sufficient to adjust the pH to a value of from 4 to 8.

Example 4

This Example also illustrates a hair tonic which is suitable for application to hair or scalp.
The hair tonic has the following formulation:

EP 0 375 388 B1

	% w/v
glucaric acid	20
ethanol	40
triethanolamine	q.s.*
perfume	q.s.
water	to 100

*sufficient to adjust to pH to a value of from 4 to 8.

Examples 5 to 8

The following formulations represent lotions which can be used topically in the treatment of bald or balding male or female heads.

	% w/v			
	5	6	7	8
Hydroxyethyl cellulose	0.4	-	0.4	-
Absolute ethanol	25	25	25	25
Propane-1,2-diol	-	-	38.4	38.4
Butane-1,3-diol	38.4	38.8	-	-
Paramethyl benzoate	0.2	0.2	0.2	0.2
Sodium talosaccharate	5	-	-	-
Sodium glucarate	-	1	-	-
di-(diethanolamine) glucarate	-	-	0.8	-
di-(triethanolamine) glucarate	-	-	-	0.6
Perfume	1	1	1	1
Water	to 100	100	100	100

Examples 9 to 12

The following formulations represent creams which can be used in the treatment of baldness.

	% w/v			
	9	10	11	12
Cetyl alcohol polyoxyethylene (10)	4	4	4	4
Cetyl alcohol	4	4	4	4
Mineral oil	4	2	-	-
Paraffin wax	-	2	4	-
Partial glyceride of palmitic and stearic acids	-	-	-	4
Glucaric acid	2	-	-	-
Monomethyl gluconate	-	-	-	1
Diethyl glucarate	-	1.5	-	-
Monooctyl glucarate	-	-	2	-
Triethanolamine	0.75	0.75	0.75	0.75
Butane-1,3-diol	3	3	3	3
Xanthan gum	0.3	0.3	0.3	0.3
Preservative	0.4	0.4	0.4	0.4
Perfume	q.s.	q.s.	q.s.	q.s.
Water	to 100	100	100	100

EP 0 375 388 B1

The following examples 13 to 17 illustrate shampoos for use in washing the hair and scalp, and for promoting hair growth on the scalp.

Example 13

	% w/v
Sodium lauryl ether sulphate (2 EO) [21% AD]	41.4
Lauryl dimethylamino acetic acid betaine: [30% AD]	4
Coconut fatty acid diethanolamine	1.5
Oleyl triethoxy phosphate (BRIPHOS 03D)	1
Polyglycol-polyamine condensation resin (POLYQUART H) [50% active]	1.5
Preservative, colouring matter, salt	0.58
Mannosaccharic acid, lactyl ester	5
Perfume	q.s.
Water	to 100

Example 14

	% w/v
Sodium lauryl ether sulphate (2 EO) [100% AD]	12
POLYMER JR400 TM	2.5
BRIPHOS TM 03D	2.5
Galactosaccharic acid, ester	4
Perfume	q.s.
Water	to 100

Example 15

	% w/v
Monoethanolamine lauryl sulphate : [100% AD]	20
JAGUAR TM C13S	3
BRIPHOS 03D	1.7
Coconut diethanolamide	5
D-Glucaro-1,4-lactone	1
Sodium allosaccharate	3
Perfume	q.s.
Water	to 100
pH adjusted to 6.5	

Example 16

	% w/v
Sodium lauryl ether sulphate (3 EO) : [100% AD]	12
JAGUAR C13S	0.3
BRIPHOS O3D	1
N-Acetylglucosamine	2
Mono (triethanolamine) altrosaccharate	4
Perfume	q.s.
Water	to 100
pH adjusted to 6.5	

Example 17

	% w/v
Sodium lauryl ether sulphate (2 EO) [100% AD]	12
POLYMER JR400	3
BRIPHOS O3D	1
Opacifier	9
Monopotassium gulosaccharate	5
Perfume	q.s.
Water	to 100
pH adjusted to 6.5	

Example 18

The following example illustrates a lotion according to the invention which can be applied topically to the scalp to prevent hair loss and stimulate hair regrowth.

	% w/v
Sodium idosaccharate	7
Minoxidil	2
ethanol	16
citric acid	1.05
water	to 100
pH adjusted to 4.2 with sodium hydroxide	

Example 19

This example illustrates a lotion for topical application to the skin in order particularly to demonstrate significant hair growth.

A concentrate was prepared by mixing together the following ingredients:

Glucaric acid solution*	30 ml
Ethanol	20 ml
10% w/w aqueous solution of TRITON-X100	1 ml
Citric acid	1 g
Water	20 ml

*The glucaric acid solution was a boiled equilibrium mixture containing 37% solids w/v, and had a pH value of 1.5. The composition of this solution, with respect to the solids was 30 parts by weight glucaro-1,4-lactone, 30 parts by weight glucaro-3,6-lactone and 40 parts by weight of glucaric acid.


The mixture was then titrated with 10N sodium hydroxide to a pH value of 4.2 to provide the concentrate.

The concentrate was finally diluted to 100 ml with water to provide the lotion.

Topical application to the rat model of the products described in Examples 1 to 19, over a period of no more than 3 months, will increase hair growth by an amount which is at least 10% more than that achievable following topical application of a similar product from which the hexosaccharic acid, esters or salts thereof have been omitted, in accordance with the Rat Hair Growth Test as described herein.

Claims

1. A preserved composition suitable for topical application to mammalian skin or hair for inducing, maintaining or increasing hair growth which comprises, as a hair growth promoter, a hexosaccharic acid and/or a salt or ester thereof; the total amount of hexosaccharic acid or salt or ester thereof present in the composition being sufficient to increase hair growth in the rat, when the composition is applied topically thereto over a period of no more than 3 months, by at least 10% more than that obtainable using a control composition from which the promoter has been omitted, in accordance with the Rat Hair Growth Test.
2. A composition according to claim 1, in which the hexosaccharic acid and/or a salt or ester thereof has the structure (1):



where

- X^1 is chosen from H, alkali metal, ammonium and substituted ammonium cations;
- X^2 is chosen from an alkyl or hydroxyalkyl group having from 1 to 18 carbon atoms;
- $\text{Y}^1, \text{Y}^2, \text{Y}^3$ and Y^4 are each chosen from H, an alkyl group having from 1 to 12 carbon atoms, and an acyl group having from 1 to 18 carbon atoms;
- l is an integer of from 1 to 3;
- m and n are each 0 or the integer 1 or 2; and
- $m+n$ is 1 or 2.


3. A composition according to claim 1 or 2, in which the hexosaccharic acid is chosen from
Allosaccharic acid
Altrosaccharic acid
Glucosaccharic acid
5 Mannosaccharic acid
Gulosaccharic acid
Idosaccharic acid
Galactosaccharic acid, and
Talosesaccharic acid.
10
4. A composition according to claim 1, 2 or 3 in which the salt of the hexosaccharic acid is a monovalent alkali metal salt, the counterion being chosen from sodium and potassium.
5. A composition according to claim 4, in which the salt of the hexosaccharic acid is the disodium salt of
15 glucosaccharic acid.
6. A composition according to claim 1, 2 or 3, in which the salt of the hexosaccharic acid is a substituted ammonium salt, the counterion being chosen from diethanolammonium and triethanolammonium.
- 20 7. A composition according to claim 1, 2 or 3, in which the ester of the hexosaccharic acid is chosen from alkyl and hydroxyalkyl esters in which the ester group has from 1 to 18 carbon atoms.
8. A composition according to any of claims 2 to 7, in which one or more of the Y¹, Y², Y³ and Y⁴ groups in structure (1) are chosen from alkyl groups having from 1 to 12 carbon atoms.
- 25 9. A composition according to any of claims 2 to 7, in which one or more of the Y¹, Y², Y³ and Y⁴ groups in structure (1) are chosen from acyl groups having from 1 to 18 carbon atoms.
10. A composition according to any preceding claim, in which the sufficient amount of the hair growth promoter present in the composition is from 0.0001 to 99% by weight.
30
11. A composition according to any preceding claim, in which the sufficient amount of the hair growth promoter present in the composition is from 1 to 50% by weight.
- 35 12. A composition according to any preceding claim, in which the amount of the hair growth promoter present in the composition is sufficient to increase hair growth in the rat by at least 20%.
13. A composition according to any preceding claim, which further comprises from 1 to 99.99% by weight of a cosmetically acceptable vehicle.
40
14. A composition according to any preceding claim, which further comprises from 0.01 to 10% by weight of a perfume.
15. A composition according to any preceding claim, which further comprises an activity enhancer.
45
16. A composition according to claim 15, in which the activity enhancer is a hair growth stimulant.
17. A composition according to claim 16, in which the hair growth stimulant is minoxidil.
- 50 18. A cosmetic method for converting vellus hair to growth as terminal hair which comprises the step of applying to the scalp in the region of vellus hair an effective amount of the composition according to any preceding claim.
- 55 19. A cosmetic method for increasing the rate of terminal hair growth, which comprises the step of applying to the scalp in the region of terminal hair an effective amount of the composition according to any of claims 1 to 17.

20. The use of the composition according to any of claims 1 to 17 for inducing, maintaining or increasing hair growth following topical application to the skin.

Patentansprüche

1. Konservierte Zusammensetzung mit Eignung zur topischen Applikation auf Säugetierhaut oder -haar zum Induzieren, Beibehalten oder Erhöhen eines Haarwachstums, umfassend als einen Haarwachstumsbeschleuniger eine Hexosaccharinsäure und/oder ein(en) Salz oder Ester davon, wobei die in der Zusammensetzung vorliegende Gesamtmenge an der Hexosaccharinsäure oder seinem Salz oder Ester ausreicht, um ein Haarwachstum bei der Ratte bei topischer Applikation der Zusammensetzung darauf über einen Zeitraum von nicht länger als drei Monaten um mindestens 10% über dem, das unter Verwendung einer Vergleichszusammensetzung ohne den Beschleuniger erhalten werden kann, gemäß der Rattenhaarwachstumsuntersuchung zu erhöhen.

2. Zusammensetzung nach Anspruch 1, bei der die Hexosaccharinsäure und/oder ihr Salz oder Ester die Struktur (1) aufweist:



worin bedeuten:

X¹ H, ein Alkalimetall, Ammonium oder ein substituiertes Ammoniumgegenion,

X² eine Alkyl- oder Hydroxyalkylgruppe mit 1 bis 18 Kohlenstoffatom(en),

Y¹, Y², Y³ und Y⁴ jeweils H, eine Alkylgruppe mit 1 bis 12 Kohlenstoffatom(en) oder eine Acylgruppe

mit 1 bis 18 Kohlenstoffatom(en),

l eine ganze Zahl von 1 bis 3,

m und n jeweils 0, 1 oder 2 und


m + n 1 oder 2.

3. Zusammensetzung nach Anspruch 1 oder 2, bei der die Hexosaccharinsäure aus Allosaccharinsäure, Altrosaccharinsäure, Glucosaccharinsäure, Mannosaccharinsäure, Gulosaccharinsäure, Idosaccharinsäure, Galactosaccharinsäure und Talosaccharinsäure ausgewählt ist.
4. Zusammensetzung nach Anspruch 1, 2 oder 3, bei der das Salz der Hexosaccharinsäure aus einem einwertigen Alkalimetallsalz, wobei das Gegenion aus Natrium und Kalium ausgewählt ist, besteht.
5. Zusammensetzung nach Anspruch 4, bei der das Salz der Hexosaccharinsäure aus dem Dinatriumsalz von Glucosaccharinsäure besteht.
6. Zusammensetzung nach Anspruch 1, 2 oder 3, bei der das Salz der Hexosaccharinsäure aus einem substituierten Ammoniumsalz, wobei das Gegenion aus Diethanolammonium und Triethanolammonium ausgewählt ist, besteht.
7. Zusammensetzung nach Anspruch 1, 2 oder 3, bei der der Ester der Hexosaccharinsäure aus Alkyl- und Hydroxyalkylester, bei denen die Estergruppe 1 bis 18 Kohlenstoffatom(e) aufweist, ausgewählt ist.
8. Zusammensetzung nach einem der Ansprüche 2 bis 7, bei der eine oder mehrere der Gruppen Y¹, Y², Y³ und Y⁴ in der Struktur (1) aus Alkylgruppen mit 1 bis 12 Kohlenstoffatom(en) ausgewählt ist.

9. Zusammensetzung nach einem der Ansprüche 2 bis 7, bei der eine oder mehr der Gruppen Y¹, Y², Y³ und Y⁴ in der Struktur (1) aus Acylgruppen mit 1 bis 18 Kohlenstoffatom(en) ausgewählt ist.
10. Zusammensetzung nach einem der vorhergehenden Ansprüche, bei der die in der Zusammensetzung vorliegende ausreichende Menge an dem Haarwachstumsbeschleuniger zwischen 0,0001 und 99 Gew.-% liegt.
11. Zusammensetzung nach einem der vorhergehenden Ansprüche, bei der die in der Zusammensetzung vorliegende ausreichende Menge an dem Haarwachstumsbeschleuniger zwischen 1 und 50 Gew.-% liegt.
12. Zusammensetzung nach einem der vorhergehenden Ansprüche, bei der die in der Zusammensetzung vorhandene Menge an dem Haarwachstumsbeschleuniger ausreicht, um ein Haarwachstum bei der Ratte um mindestens 20% zu erhöhen.
13. Zusammensetzung nach einem der vorhergehenden Ansprüche, die des weiteren 1 bis 99,99 Gew.-% eines kosmetisch akzeptablen Trägers umfaßt.
14. Zusammensetzung nach einem der vorhergehenden Ansprüche, die des weiteren 0,01 bis 10 Gew.-% eines Duftstoffs umfaßt.
15. Zusammensetzung nach einem der vorhergehenden Ansprüche, die des weiteren ein eine Aktivität erhöhendes Mittel umfaßt.
16. Zusammensetzung nach Anspruch 15, bei der das eine Aktivität erhöhende Mittel aus einem ein Haarwachstum stimulierenden Mittel besteht.
17. Zusammensetzung nach Anspruch 16, bei der das ein Haarwachstum stimulierende Mittel Minoxidil ist.
18. Ein kosmetisches Verfahren zur Umwandlung von Flaumhaar in ein Wachstum als Terminalhaar, umfassend den Schritt Applizieren einer wirksamen Menge der Zusammensetzung gemäß einem der vorhergehenden Ansprüche auf die Kopfhaut im Bereich von Flaumhaar.
19. Kosmetisches Verfahren zur Erhöhung der Rate an Terminalhaarwachstum, umfassend die Stufe Applizieren einer wirksamen Menge der Zusammensetzung gemäß einem der Ansprüche 1 bis 17 auf die Kopfhaut im Bereich von Terminalhaar.
20. Verwendung der Zusammensetzung gemäß einem der Ansprüche 1 bis 17 zum Induzieren, Beibehalten oder Erhöhen eines Haarwachstums nach einer topischen Applikation auf die Haut.

Revendications

1. Une composition conservée appropriée pour être appliquée de façon topique sur la peau ou sur les cheveux ou les poils des mammifères pour provoquer, conserver ou augmenter la pousse des cheveux ou des poils, comprenant en tant que promoteur de pousse des cheveux ou des poils, un acide hexosaccharique et/ou un sel ou un ester de celui-ci ; la quantité totale d'acide hexosaccharique ou de sel ou d'ester de celui-ci présente dans la composition étant suffisante pour augmenter la pousse des poils chez le rat, lorsque la composition y est appliquée de façon topique sur une période n'excédant pas 3 mois, d'au moins 10% de plus que ce qu'on peut obtenir en utilisant une composition témoin ne contenant pas de promoteur, en accord avec le Test de Pousse des Poils chez le Rat.
2. Une composition selon la revendication 1, dans laquelle l'acide hexosaccharique et/ou un sel ou un ester de celui-ci a la structure (1) :



où

X¹ est de l'hydrogène, un métal alcalin, ou un ion antagoniste ammonium ou ammonium substitué;
X² est choisi à partir d'un groupe alkyle ou hydroxyalkyle ayant de 1 à 18 atomes de carbone;
Y¹, Y², Y³ et Y⁴ sont chacun choisis à partir de H, d'un groupe alkyle ayant de 1 à 12 atomes de carbone, et d'un groupe acyle ayant de 1 à 18 atomes de carbone;

1 est un nombre entier allant de 1 à 3;

m et n sont chacun 0 ou l'entier 1 ou 2 ; et

$m + n$ est 1 ou 2.

- 3. Une composition selon la revendication 1 ou 2, dans laquelle l'acide hexosaccharique est choisi parmi :**

l'acide allosaccharique

l'acide altrosaccharique

l'acide glucosaccharique

l'acide mannosaccharique

l'acide gulosaccharique

l'acide idosaccharique

l'acide galactosaccharique, et

l'acide talosaccharique.

4. Une composition selon la revendication 1, 2 ou 3, dans laquelle le sel de l'acide hexosaccharique est un sel de métal alcalin monovalent, l'ion antagoniste étant choisi parmi le sodium et le potassium.

5. Une composition selon la revendication 4, dans laquelle le sel d'acide hexosaccharique est le sel de disodium d'acide glucosaccharique.

6. Une composition selon la revendication 1, 2 ou 3, dans laquelle le sel de l'acide hexosaccharique est un sel d'ammonium substitué, l'ion antagoniste étant choisi à parmi le diéthanolammonium et le triéthanolammonium.

7. Une composition selon la revendication 1, 2 ou 3, dans laquelle l'ester de l'acide hexosaccharique est choisi parmi les esters alkyles et hydroxyalkyles dans lesquels le groupe ester a de 1 à 18 atomes de carbone.

8. Une composition selon l'une des revendications 2 à 7, dans laquelle un ou plusieurs des groupes Y¹, Y², Y³ et Y⁴ dans la structure (1) sont choisis parmi les groupes alkyles ayant de 1 à 12 atomes de carbone.

9. Une composition selon l'une des revendications 2 à 7, dans laquelle un ou plusieurs des groupes Y¹, Y², Y³ et Y⁴ dans la structure (1) sont choisis parmi les groupes acyles ayant de 1 à 18 atomes de carbone.

- 55 10. Une composition selon l'une des revendications précédentes, dans laquelle la quantité suffisante du promoteur de pousse des cheveux ou des poils présente dans la composition va de 0,0001 à 99% en masse.

11. Une composition selon l'une des revendications précédentes, dans laquelle la quantité suffisante du promoteur de pousse des cheveux ou des poils présente dans la composition va de 1 à 50% en masse.
- 5 12. Une composition selon l'une des revendications précédente, dans laquelle la quantité du promoteur de pousse des cheveux ou des poils présente dans la composition est suffisante pour augmenter la pousse des poils chez le rat d'au moins 20%.
- 10 13. Une composition selon l'une des revendications précédentes, comprenant en outre de 1 à 99,99% en masse d'un véhicule acceptable d'un point de vue cosmétique.
14. Une composition selon l'une des revendications précédentes, comprenant en outre de 0,01 à 10% en masse d'un parfum.
- 15 15. Une composition selon l'une des revendications précédentes, comprenant en outre un agent d'accroissement de l'activité.
16. Une composition selon la revendication 15, dans laquelle l'agent d'accroissement de l'activité est un stimulant de la pousse des cheveux.
- 20 17. Une composition selon la revendication 16, dans laquelle le stimulant de pousse des cheveux est du minoxidil.
- 25 18. Un procédé cosmétique pour transformer le duvet et le faire pousser sous forme de cheveux définitifs, comprenant l'étape consistant à appliquer sur le cuir chevelu dans la zone où il y a le duvet, une quantité efficace de la composition en accord avec l'une des revendications précédentes.
- 30 19. Un procédé cosmétique pour augmenter la vitesse de pousse des cheveux définitifs, comprenant l'étape consistant à appliquer sur le cuir chevelu au niveau de la zone des cheveux définitifs une quantité efficace de la composition selon l'une des revendications 1 à 17.
20. L'utilisation de la composition selon l'une des revendications 1 à 17 pour provoquer, conserver ou augmenter la pousse des cheveux ou des poils suite à une application topique sur la peau.

35

40

45

50

55

THIS PAGE BLANK (USPTO)